

LOCAL HOMOLOGICAL PROPERTIES AND CYCLICITY OF HOMOGENEOUS ANR COMPACTA

V. VALOV

ABSTRACT. In accordance with the Bing-Borsuk conjecture [2], we show that if X is an n -dimensional homogeneous metric ANR compactum and $x \in X$, then there is a local basis at x consisting of connected open sets U such that the homological properties of \overline{U} and bdU are similar to the properties of the closed ball $\mathbb{B}^n \subset \mathbb{R}^n$ and its boundary \mathbb{S}^{n-1} . We discuss also the following questions raised by Bing-Borsuk [2], where X is a homogeneous ANR -compactum with $\dim X = n$:

- Is it true that X is cyclic in dimension n ?
- Is it true that no non-empty closed subset of X , acyclic in dimension $n - 1$, separates X ?

It is shown that both questions have positive answers simultaneously, and a positive solution to each one of them implies a solution to another question of Bing-Borsuk (whether every finite-dimensional homogenous metric AR -compactum is a point).

1. INTRODUCTION

There are few open problems concerning homogeneous compacta, see [2]. The most important one is the well known Bing-Borsuk conjecture stating that every n -dimensional homogeneous metric ANR -compactum X is an n -manifold. Another one is whether any such X has the following properties: (i) X is cyclic in dimension n ; (ii) no closed non-empty subset of X , acyclic in dimension $n - 1$, separates X . It is also unknown if there exists a non-trivial finite dimensional metric homogeneous AR compactum.

In this paper we address the above problems and investigate the homological structure of homogeneous metric ANR -compacta. In accordance with the Bing-Borsuk conjecture, we prove that any such a

2010 *Mathematics Subject Classification*. Primary 55M10, 55M15; Secondary 54F45, 54C55.

Key words and phrases. Bing-Borsuk conjecture for homogeneous compacta, dimensionally full-valued compacta, homology membrane, homological dimension, homology groups, homogeneous metric ANR -compacta.

The author was partially supported by NSERC Grant 261914-13.

compactum has local homological properties similar to the local structure of \mathbb{R}^n , see Theorem 1.1. It is also shown that the properties (i) and (ii) from the second of the above questions are equivalent, so each one of them implies that every finite-dimensional homogeneous metric ANR is a point.

Reduced Čech homology $H_n(X; G)$ and cohomology groups $H^n(X; G)$ with coefficient from G are considered everywhere below, where G is an abelian group. Suppose (K, A) is a pair of closed subsets of a space X with $A \subset K$. By $i_{A,K}^n$ we denote the homomorphism from $H_n(A; G)$ into $H_n(K; G)$ generated by the inclusion $A \hookrightarrow K$. Following [2], we say that K is an n -homology membrane spanned on A for an element $\gamma \in H_n(A; G)$ provided γ is homologous to zero in K , but not homologous to zero in any proper closed subset of K containing A . It is well known [2, Property 5., p.103] that for every compact metric space X and a closed set $A \subset X$ the existence a non-trivial element $\gamma \in H_n(A; G)$ with $i_{A,X}^n(\gamma) = 0$ yields the existence of a closed set $K \subset X$ containing A such that K is an n -homology membrane for γ spanned on A . We also say that a space K is a homological (n, G) -bubble if $H_n(K; G) \neq 0$, but $H_n(B; G) = 0$ for every closed proper subset $B \subset K$.

For any abelian group G Alexandroff [1] introduced the dimension $d_G X$ of a space X as the maximum integer n such that there exist a closed set $F \subset X$ and a nontrivial element $\gamma \in H_{n-1}(F; G)$ with γ being G -homologous to zero in X . According to [1, p.207] we have the following inequalities for any metric finite-dimensional compactum X : $d_G X \leq \dim X$ and $\dim X = d_{\mathbb{Q}_1} X = d_{\mathbb{S}^1} X$, where G is any abelian group, \mathbb{S}^1 is the circle group and \mathbb{Q}_1 is the group of rational elements of \mathbb{S}^1 .

Because the definition of $d_G X$ does not provide any information for the homology groups $H_{k-1}(F; G)$ when $F \subset X$ is closed and $k < d_G X - 1$, we consider the set $\mathcal{H}_{X,G}$ of all integers $k \geq 1$ such that there exist a closed set $F \subset X$ and a nontrivial element $\gamma \in H_{k-1}(F; G)$ with $i_{F,X}^{k-1}(\gamma) = 0$. Obviously, $d_G X = \max \mathcal{H}_{X,G}$.

Using the properties of the sets $\mathcal{H}_{X,G}$, we investigate in Section 2 the local homological properties of metric homogeneous ANR compacta. The main result in that section is Theorem 1.1 below, which is a homological version of [11, Theorem 1.1].

Theorem 1.1. *Let X be a finite dimensional homogeneous metric ANR with $\dim X \geq 2$. Then every point $x \in X$ has a basis $\mathcal{B}_x = \{U_k\}$ of open sets such that for any abelian group G and $n \geq 2$ with $n \in \mathcal{H}_{X,G}$ and $n + 1 \notin \mathcal{H}_{X,G}$ almost all U_k satisfy the following conditions:*

- (1) $H_{n-1}(\text{bd}\overline{U}_k; G) \neq 0$ and \overline{U}_k is an $(n-1)$ -homology membrane spanned on $\text{bd}\overline{U}_k$ for any non-zero $\gamma \in H_{n-1}(\text{bd}\overline{U}_k; G)$;
- (2) $H_{n-1}(\overline{U}_k; G) = H_n(\overline{U}_k; G) = 0$ and $X \setminus \overline{U}_k$ is connected;
- (3) $\text{bd}\overline{U}_k$ is a homological $(n-1, G)$ -bubble.

Corollary 1.2. *Let X be as in Theorem 1.1. Then X has the following property for any abelian group G and $n \geq 2$ with $n \in \mathcal{H}_{X,G}$ and $n+1 \notin \mathcal{H}_{X,G}$: If a closed subset $K \subset X$ is an $(n-1)$ -homology membrane spanned on B for some closed set $B \subset X$ and $\gamma \in H_{n-1}(B; G)$, then $(K \setminus B) \cap \overline{X \setminus K} = \emptyset$.*

In Section 3 we show that the following two statements are equivalent, where $\mathcal{H}(n)$ is the class of all homogeneous metric ANR-compacta X with $\dim X = n$:

- (1) For all $n \geq 1$ and $X \in \mathcal{H}(n)$ there is a group G such that $H^n(X; G) \neq 0$ (resp., $H_n(X; G) \neq 0$);
- (2) If $X \in \mathcal{H}(n)$ with $n \geq 1$, and $F \subset X$ is a closed separator of X with $\dim F = n-1$, then there exists a group G such that $H^{n-1}(F; G) \neq 0$ (resp., $H_{n-1}(F; G) \neq 0$).

Therefore, we have the following result (see Corollary 3.3):

Theorem 1.3. *Suppose for all $n \geq 1$ and all $X \in \mathcal{H}(n)$ the following holds: For every closed separator F of X with $\dim F = n-1$ there exists a group G such that either $H^{n-1}(F; G) \neq 0$ or $H_{n-1}(F; G) \neq 0$. Then there is no homogeneous metric AR-compactum Y with $\dim Y < \infty$.*

2. LOCAL HOMOLOGICAL PROPERTIES OF HOMOGENEOUS ANR-COMPACTA

We begin this section with the following analogue of Theorem 8.1 from [2].

Proposition 2.1. *Let X be a locally compact and homogeneous separable metrizable ANR space. Suppose there is a pair $F \subset K$ of compact proper subsets of X such that K is contractible in X and K is a homological membrane for some $\gamma \in H_{n-1}(F; G)$. If $(K \setminus F) \cap \overline{X \setminus K} \neq \emptyset$, then there exists of a proper compact subset $P \subset X$ contractible in X such that $H_n(P; G) \neq 0$.*

Proof. We follow the proof of [6, Lemma 1] (let us note that the proof of Proposition 3.1 can also be obtained following the arguments of [2, Theorem 8.1]). Let $a \in (K \setminus F) \cap \overline{X \setminus K}$. Then a is a boundary point for K . Because K is contractible in X , there is a homotopy $g : K \times [0, 1] \rightarrow X$ such that $g(x, 0) = x$ and $g(x, 1) = c \in X$ for

all $x \in K$. Then we can find an open set $U \subset X$ containing K and a homotopy $\bar{g} : \bar{U} \times [0, 1] \rightarrow X$ extending g and connecting the identity on \bar{U} and the constant map $\bar{U} \rightarrow c$ (this can be done since X is an ANR). So, \bar{U} is also contractible in X . Moreover, we can assume that \bar{U} is compact. Fix a metric d on X generating its topology in the following way: consider X as a subspace of its one-point compactification αX and take d to be the restriction to X of some admissible metric on αX . Let $2\epsilon = d(a, F)$ and take an open cover ω of U such that for any two ω -close maps $f_1, f_2 : K \rightarrow U$ (i.e., for all $x \in K$ the points $f_1(x), f_2(x)$ are contained in some element of ω) there is an ϵ -homotopy $\Phi : K \times [0, 1] \rightarrow U$ between f_1 and f_2 (i.e., each set $M_\Phi(x) = \{\Phi(x, t) : t \in [0, 1]\}$, $x \in K$, is of diameter $< \epsilon$). This can be done because U is an ANR . Now, we fix an open set $V \subset X$ containing K with $\bar{V} \subset U$ and let δ be the Lebesgue's number of the open cover $\{\Gamma \cap \bar{V} : \Gamma \in \omega\}$ of \bar{V} . According to Effros' theorem [4], there is a positive number η such that if $x, y \in X$ are two points with $d(x, y) < \eta$, then $f(x) = y$ for some homeomorphism $h : X \rightarrow X$, which is $\min\{\delta, d(K, X \setminus V)\}$ -close to the identity on X (the Effros' theorem can be applied because of the special choice of the metric d). Since a is a boundary point for K , we can choose a point $b \in V \setminus K$ with $d(a, b) < \eta$. Then, there exists a homeomorphism $h'_1 : X \rightarrow X$ such that $h'_1(a) = b$ and $d(x, h'_1(x)) < \min\{\delta, d(K, X \setminus V)\}$, $x \in X$. Let h_1 be the restriction $h'_1|_K$. Obviously, $h_1 : K \rightarrow h_1(K)$ is a homeomorphism with $h_1(K) \subset V$ and h_1 is δ -close to the identity on K . Then, according to the choice of δ , there is homotopy $h : K \times [0, 1] \rightarrow U$ such that $h(x, 0) = x$, $h(x, 1) = h_1(x)$ and $d(x, h(x, t)) < \epsilon$ for all $x \in K$ and $t \in [0, 1]$.

Let $K_1 = K \cup h(F \times \mathbb{I})$, $K_2 = h_1(K)$ and $K_0 = K_1 \cap K_2$, where $\mathbb{I} = [0, 1]$. Since $2\epsilon = d(a, F)$ and h is an ϵ -small homotopy, $b \in K_2 \setminus K_1$. So, K_0 is a proper subset of K_2 containing $h_1(F)$. Hence, $D = h_1^{-1}(K_0)$ is a proper subset of K containing F , which implies $\gamma_1 = i_{F,D}^{n-1}(\gamma) \neq 0$. Because h_1 is a homeomorphism, $(\varphi_1)_* : H_{n-1}(D; G) \rightarrow H_{n-1}(K_0; G)$ is an isomorphism, where $\varphi_1 = h_1|_D$. Thus, $\hat{\gamma} = (\varphi_1)_*(\gamma_1) \neq 0$.

Claim 1. $i_{K_0, K_1}^{n-1}(\hat{\gamma}) = 0$ and $i_{K_0, K_2}^{n-1}(\hat{\gamma}) = 0$.

Let $\lambda = i_{F, h(F \times \mathbb{I})}^{n-1}(\gamma)$. Since $h|(F \times \mathbb{I})$ is a homotopy between the identity on F and the map $\varphi_2 = h_1|_F$, $\lambda = i_{h_1(F), h(F \times \mathbb{I})}^{n-1}((\varphi_2)_*(\gamma))$. We consider the following commutative diagram

$$\begin{array}{ccc}
\gamma \in H_{n-1}(F; G) & \xrightarrow{i_{F,D}^{n-1}} & H_{n-1}(D; G) \ni \gamma_1 \\
(\varphi_2)_* \downarrow & & (\varphi_1)_* \downarrow \\
(\varphi_2)_*(\gamma) \in H_{n-1}(h_1(F); G) & \xrightarrow{i_{h_1(F), K_0}^{n-1}} & H_{n-1}(K_0; G) \ni \hat{\gamma} \\
i_{h_1(F), h(F \times \mathbb{I})}^{n-1} \downarrow & & \nearrow i_{K_0, K_1}^{n-1} \\
\lambda \in H_{n-1}(h(F \times \mathbb{I}); G) & & \\
i_{h(F \times \mathbb{I}), K_1}^{n-1} \downarrow & & \\
0 \in H_{n-1}(K_1; G) & &
\end{array}$$

Obviously, $i_{F, K_1}^{n-1}(\gamma) = i_{h(F \times \mathbb{I}), K_1}^{n-1}(\lambda) = i_{K_0, K_1}^{n-1}(\hat{\gamma})$. On the other hand, $i_{F, K_1}^{n-1}(\gamma) = i_{K, K_1}^{n-1}(i_{F, K}^{n-1}(\gamma)) = 0$ because $i_{F, K}^{n-1}(\gamma) = 0$. Hence, $i_{K_0, K_1}^{n-1}(\hat{\gamma}) = 0$.

For the second part of the claim, observe that $i_{D, K}^{n-1}(\gamma_1) = i_{F, K}^{n-1}(\gamma) = 0$. Then, the equality $i_{K_0, K_2}^{n-1}(\hat{\gamma}) = 0$ follows from next diagram

$$\begin{array}{ccc}
\gamma_1 \in H_{n-1}(D; G) & \xrightarrow{i_{D, K}^{n-1}} & H_{n-1}(K; G) \ni 0 \\
(\varphi_1)_* \downarrow & & (h_1)_* \downarrow \\
\hat{\gamma} \in H_{n-1}(K_0; G) & \xrightarrow{i_{K_0, K_2}^{n-1}} & H_{n-1}(K_2; G) \ni 0
\end{array}$$

We are in a position now to complete the proof of Proposition 3.1. Let $P = K_1 \cup K_2$. Since $h(K \times \mathbb{I}) \subset U$, $P \subset U$. Therefore, P is contractible in X (recall that \bar{U} is contractible in X). Finally, Claim 1 and the Mayer-Vietoris exact sequence

$$H_n(P; G) \xrightarrow{\partial} H_{n-1}(K_0; G) \xrightarrow{\psi} H_{n-1}(K_1; G) \oplus H_{n-1}(K_2; G),$$

where $\psi = i_{K_0, K_1}^{n-1} \oplus i_{K_0, K_2}^{n-1}$, show that there exists a non-trivial $\alpha \in H_n(P; G)$ with $\partial(\alpha) = (\hat{\gamma})$. \square

For simplicity, we say that a closed set $F \subset X$ is *strongly contractible* in X if F is contractible in a closed set $A \subset X$ and A is contractible in X .

Corollary 2.2. *Let X be a homogeneous compact metrizable ANR-space such that $n \in \mathcal{H}_{X, G}$ and $n+1 \notin \mathcal{H}_{X, G}$. Then for every closed set $F \subset X$ we have:*

- (1) $H_n(F; G) = 0$ provided F is contractible in X ;
- (2) F separates X provided $H_{n-1}(F; G) \neq 0$ and F is strongly contractible in X ;

- (3) If K is a homological membrane for some non-trivial element of $H_{n-1}(F; G)$ and K is contractible in X , then $(K \setminus F) \cap \overline{X \setminus K} = \emptyset$.

Proof. Since F is contractible in X , every $\gamma \in H_n(F; G)$ is homologous to zero in X . So, the existence of a non-trivial element of $H_n(F; G)$ would imply $n + 1 \in \mathcal{H}_{X,G}$, a contradiction.

To prove the second item, suppose $H_{n-1}(F; G) \neq 0$ and F is strongly contractible in X . So, there exists a closed set $A \subset X$ such that F is contractible in A and A is contractible in X . Then, by [2, Property 5., p.103], we can find a closed set $K \subset A$ containing F which is a homological membrane for some non-trivial $\gamma \in H_{n-1}(F; G)$. Because K (as a subset of A) is contractible in X , the assumption $(K \setminus F) \cap \overline{X \setminus K} \neq \emptyset$ would yield the existence of a proper closed set $P \subset X$ contractible in X with $H_n(P; G) \neq 0$ (see Proposition 2.1). Consequently, there would be a non-trivial $\alpha \in H_n(P; G)$ homologous to zero in X . Hence, $n + 1 \in \mathcal{H}_{X,G}$, a contradiction. Therefore, $(K \setminus F) \cap \overline{X \setminus K} = \emptyset$. This means that $X \setminus F = (K \setminus F) \cup (X \setminus K)$ with both $K \setminus F$ and $X \setminus K$ being non-empty open disjoint subsets of X .

The above arguments provide also the proof of the third item. \square

Proof of Theorem 1.1. Suppose X satisfies the hypotheses of Theorem 1.1. By [11, Theorem 1.1], every $x \in X$ has a basis $\mathcal{B}_x = \{U_k\}_{k \geq 1}$ of open sets satisfying the following conditions: $\text{bd}U_k = \text{bd}\overline{U}_k$; the sets U_k , $\text{bd}\overline{U}_k$ and $X \setminus \overline{U}_k$ are connected; $H^{\dim X - 1}(A; \mathbb{Z}) = 0$ for all proper closed sets $A \subset \text{bd}\overline{U}_k$. We may also suppose that each \overline{U}_{k+1} is contractible in U_k and all \overline{U}_k are strongly contractible in X . Let G be an abelian group and $n \geq 2$ with $n + 1 \notin \mathcal{H}_{X,G}$ and $n \in \mathcal{H}_{X,G}$. So, there exist a closed set $B \subset X$ and a nontrivial element $\gamma \in H_{n-1}(B; G)$ with $i_{F,X}^{n-1}(\gamma) = 0$. Then, by [2, Property 5., p.103], there is a closed set $K \subset X$ containing B which is a homological membrane for γ . We fix a point $\tilde{x} \in K \setminus B$ and its open in K neighborhood W with $\overline{W} \cap B = \emptyset$. According to [2, Property 6., p.103], \overline{W} is an $(n-1)$ -homological membrane for some non-trivial element of $H_{n-1}(\text{bd}_K \overline{W}; G)$. We can choose W so small that \overline{W} is contractible in X . Then Corollary 2.2 yields $(\overline{W} \setminus \text{bd}_K \overline{W}) \cap \overline{X \setminus \overline{W}} = \emptyset$. So, $\overline{W} \setminus \text{bd}_K \overline{W}$ is open in X and contains \tilde{x} . Hence, there exists k_0 such that $U_k \subset \overline{W} \setminus \text{bd}_K \overline{W}$ for all $U_k \in \mathcal{B}_{\tilde{x}}$ with $k \geq k_0$. Below we consider only the elements U_k with $k \geq k_0$. Applying again [2, Property 6., p.103], we conclude that every \overline{U}_k is a homological membrane for some non-trivial element of $H_{n-1}(\text{bd}U_k; G)$. By Corollary 2.2(1), $H_n(\overline{U}_k; G) = 0$. Suppose $\gamma \in H_{n-1}(\text{bd}U_k; G)$ is non-trivial. Since $X \setminus \overline{U}_k$ is connected, Corollary 2.2(2) implies

that $H_{n-1}(\overline{U}_k; G) = 0$. Consequently, γ is homologous to zero in \overline{U}_k . So, by [2, Property 5., p.103], \overline{U}_k contains a closed set P such that P is a homological membrane for γ . Then Corollary 2.2(3) implies $(P \setminus \text{bd}U_k) \cap \overline{X \setminus P} = \emptyset$. Hence, $X \setminus \text{bd}U_k$ is the union of the disjoint open sets $P \setminus \text{bd}U_k$ and $X \setminus P$. Because U_k is connected and $U_k \cap P \neq \emptyset$, $U_k \subset P \setminus \text{bd}U_k$. Therefore, $P = \overline{U}_k$. This provides the proof of the first two conditions of Theorem 1.1.

To prove the last item of Theorem 1.1, assume that $H_{n-1}(F; G) \neq 0$ for some closed proper subset F of $\text{bd}U_{k+1}$, where $k \geq k_0$. Because F (as a subset of \overline{U}_{k+1}) is strongly contractible in X , according to Corollary 2.2(2), F separates X . So, $X \setminus F$ is the union of two disjoint non-empty open in X sets V_1 and V_2 with $\overline{V_1} \cap \overline{V_2} \subset F$. Let us show that F separates \overline{U}_k . Indeed, otherwise $\overline{U}_k \setminus F$ would be connected. Then $\overline{U}_k \setminus F$ should be contained in one of the sets V_1, V_2 , say V_1 . Since $X \setminus \overline{U}_k$ is also connected and $V_2 \neq \emptyset$, $X \setminus \overline{U}_k \subset V_2$. Hence, $\overline{U}_k \setminus F \cap X \setminus \overline{U}_k \subset F$. On the other hand, because $\overline{U}_k \setminus F$ is dense in \overline{U}_k (recall that F does not contain interior points), $\overline{U}_k \setminus F \cap X \setminus \overline{U}_k = \text{bd}U_k$. So, $F \supset \text{bd}U_k$, a contradiction. Therefore, F separates \overline{U}_k .

The proof of Theorem 1.1(3) will be done if we show that F can not separate \overline{U}_k . According to [11, Theorem 1.1], \overline{U}_k is an $(m-1)$ -cohomology membrane spanned on $\text{bd}U_k$ for some non-trivial $\alpha \in H^{m-1}(\text{bd}U_k; \mathbb{Z})$, where $m = \dim X$. This means that α (considered as a map from $\text{bd}U_k$ to the Eilenberg-MacLane complex $K(\mathbb{Z}, m-1)$) is not extendable over \overline{U}_k but it is extendable over any proper closed subset of \overline{U}_k . Hence, by [9, Proposition 2.10], the couple $(\overline{U}_k, \text{bd}U_k)$ is a strong $K_{\mathbb{Z}}^m$ -manifold (see [9] for the definition of a strong $K_{\mathbb{Z}}^m$ -manifold). So, according to [9, Theorem 3.3], $H^{m-1}(F; \mathbb{Z}) \neq 0$ because F separates \overline{U}_k and $F \cap \text{bd}U_k = \emptyset$. Finally, we obtained a contradiction because $H^{m-1}(A; \mathbb{Z}) = 0$ for every proper closed subset A of $\text{bd}U_{k+1}$. Therefore, the all U_k , $k \geq k_0 + 1$, satisfy conditions (1) – (3) from Theorem 1.1. \square

Proof of Corollary 1.2. Suppose there exists a point $a \in (K \setminus B) \cap \overline{X \setminus K}$ and take a set $U \in \mathcal{B}_a$ satisfying condition (1) – (3) from Theorem 1.1 such that $U \cap B = \emptyset$. Then $F_U = \text{bd}_K(U \cap K)$ is non-empty and it follows from [2, Property 6., p.103] that $\overline{U} \cap \overline{K}$ is a homology membrane for some non-zero $\alpha \in H_{n-1}(F_U; G)$. Because $F_U \subset \text{bd}U$, by Theorem 1.1(3), $F_U = \text{bd}U$. So, \overline{U} is a homological membrane for α , see Theorem 1.1(1). This implies that $i_{\text{bd}U, \overline{U} \cap \overline{K}}^{n-1}(\alpha) \neq 0$ provided $\overline{U} \cap \overline{K}$ is a proper subset of \overline{U} . Therefore, $\overline{U} \cap \overline{K} = \overline{U}$, which yields

$U \subset K$. The last inclusion contradicts the fact that $a \in \overline{X \setminus K}$. Hence, $(K \setminus B) \cap \overline{X \setminus K} = \emptyset$. \square

3. CYCLICITY OF HOMOGENEOUS ANR'S

Let $\mathcal{H}(n)$ be the class of all homogeneous metric ANR-compacta X with $\dim X = n$.

Theorem 3.1. *The following conditions are equivalent:*

- (1) *If $n \geq 1$, then for every space $X \in \mathcal{H}(n)$ there exists a group G with $H^n(X; G) \neq 0$;*
- (2) *If $n \geq 1$ and $X \in \mathcal{H}(n)$, then for every closed set $F \subset X$ separating X there exists a group G with $H^{n-1}(F; G) \neq 0$;*
- (3) *If $n \geq 1$ and $X \in \mathcal{H}(n)$, then for every $(n-1)$ -dimensional closed set $F \subset X$ separating X there exists a group G with $H^{n-1}(F; G) \neq 0$.*

Proof. (1) \Rightarrow (2) Suppose $n \geq 1$ and $X \in \mathcal{H}(n)$. Then $H^n(X; G) \neq 0$ for some group G , and by [10, Corollary 1.2], $H^{n-1}(F; G) \neq 0$ for every non-empty closed set $F \subset X$ separating X .

(2) \Rightarrow (3) This implication is trivial.

(3) \Rightarrow (1) Suppose that condition (3) holds, but there exists $n \geq 1$ and $X \in \mathcal{H}(n)$ such that $H^n(X; G) = 0$ for all groups G . Consider the two-dimensional sphere \mathbb{S}^2 and a circle \mathbb{S}^1 separating \mathbb{S}^2 . Then $X \times \mathbb{S}^2 \in \mathcal{H}(n+2)$ and $X \times \mathbb{S}^1$ is a closed separator of $X \times \mathbb{S}^2$ of dimension $n+1$. So, there is a group G' such that $H^{n+1}(X \times \mathbb{S}^1; G') \neq 0$. On the other hand, according to the Künneth formula, we have the exact sequence

$$\sum_{i+j=n+1} H^i(X) \otimes H^j(\mathbb{S}^1) \rightarrow H^{n+1}(X \times \mathbb{S}^1) \rightarrow \sum_{i+j=n+2} H^i(X) * H^j(\mathbb{S}^1),$$

where the coefficient group G' is suppressed. Because $\dim X = n$ and $\dim \mathbb{S}^1 = 1$, $H^{n+i}(X; G') = 0$ and $H^{1+i}(\mathbb{S}^1; G') = 0$ for all $i \geq 1$. Moreover, $H^n(X; G') = 0$. So,

$$\sum_{i+j=n+1} H^i(X; G') \otimes H^j(\mathbb{S}^1; G') = \sum_{i+j=n+2} H^i(X; G') * H^j(\mathbb{S}^1; G') = 0.$$

Hence, $H^{n+1}(X \times \mathbb{S}^1; G') = 0$, a contradiction. \square

A homological version of Theorem 3.1 also holds.

Theorem 3.2. *The following conditions are equivalent:*

- (1) *If $n \geq 1$, then for every space $X \in \mathcal{H}(n)$ there exists a group G with $H_n(X; G) \neq 0$;*

- (2) If $n \geq 1$ and $X \in \mathcal{H}(n)$, then for every closed set $F \subset X$ separating X there exists a group G with $H_{n-1}(F; G) \neq 0$;
- (3) If $n \geq 1$ and $X \in \mathcal{H}(n)$, then for every $(n-1)$ -dimensional closed set $F \subset X$ separating X there exists a group G with $H_{n-1}(F; G) \neq 0$.

Proof. Everywhere below \widehat{H}_* denotes the exact homology (see [5], [7]), which for locally compact metric spaces is equivalent to Steenrod's homology [8]. For every compact metric space X and every k there exists a surjective homomorphism $T_X^k : \widehat{H}_k(X; G) \rightarrow H_k(X; G)$. According to [7, Theorem 4 and p.125], T_X^k is an isomorphism in each of the following cases: G is a vector space over a field; both $\widehat{H}_k(X; G)$ and G are countable modules; $\dim X = k$; $H^{k+1}(X; \mathbb{Z})$ is finitely generated.

(1) \Rightarrow (2) Suppose $n \geq 1$ and $X \in \mathcal{H}(n)$. Then $H_n(X; G) \neq 0$ for some group G . By [7, Theorem 3], we have the exact sequence

$$(*) \quad \text{Ext}(H^{n+1}(X; \mathbb{Z}), G) \rightarrow \widehat{H}_n(X; G) \rightarrow \text{Hom}(H^n(X; \mathbb{Z}), G) \rightarrow 0.$$

Since $\dim X = n$, $H^{n+1}(X; \mathbb{Z}) = 0$. Moreover $\widehat{H}_n(X; G)$ is non-trivial because so is $H_n(X; G)$ and T_X^n is a surjective homomorphism. Hence, $H^n(X; \mathbb{Z}) \neq 0$ and there exists a non-trivial homomorphism $\varphi : H^n(X; \mathbb{Z}) \rightarrow G$. Now, let $F \subset X$ be a closed separator of X and $X \setminus F = X_1 \cup X_2$, where $X_1, X_2 \subset X$ are closed proper subsets with $X_1 \cap X_2 = F$. Since $H^n(P; \mathbb{Z}) = 0$ for every closed proper subset $P \subset X$ (see [10]), $H^n(F; \mathbb{Z}) = H^n(X_1; \mathbb{Z}) = H^n(X_2; \mathbb{Z}) = 0$. Then it follows from the Mayer-Vietoris sequence

$$H^{n-1}(F; \mathbb{Z}) \xrightarrow{\partial} H^n(X; \mathbb{Z}) \xrightarrow{\psi} H^n(X_1; \mathbb{Z}) \oplus H^n(X_2; \mathbb{Z})$$

that $H^{n-1}(F; \mathbb{Z}) \neq 0$ and ∂ is a surjective homomorphism. Consequently, $\varphi \circ \partial : H^{n-1}(F; \mathbb{Z}) \rightarrow G$ is also a non-trivial surjective homomorphism. Hence, $\text{Hom}(H^{n-1}(F; \mathbb{Z}), G) \neq 0$ and the exact sequence

$$0 \rightarrow \text{Ext}(H^n(F; \mathbb{Z}), G) \rightarrow \widehat{H}_{n-1}(F; G) \rightarrow \text{Hom}(H^{n-1}(F; \mathbb{Z}), G) \rightarrow 0$$

yields $\widehat{H}_{n-1}(F; G) \neq 0$. Finally, since $H^n(F; \mathbb{Z}) = 0$, $\widehat{H}_{n-1}(F; G)$ is isomorphic to $H_{n-1}(F; G)$.

(2) \Rightarrow (3) This implication is obvious.

(3) \Rightarrow (1) As in the proof of Theorem 3.1, (3) \Rightarrow (1), suppose there exists $n \geq 1$ and $X \in \mathcal{H}(n)$ such that $H_n(X; G) = 0$ for all groups G . Since $\widehat{H}_n(X; G)$ is isomorphic to $H_n(X; G)$ and $H^{n+1}(X; \mathbb{Z}) = 0$ (recall that $\dim X = n$), it follows from the exact sequence (*) that $\text{Hom}(H^n(X; \mathbb{Z}), G) = 0$ for all groups G . This implies $H^n(X; \mathbb{Z}) = 0$. As above, the product $X \times \mathbb{S}^1$ is a closed separator of $X \times \mathbb{S}^2$, and

according to our assumption, $H_{n+1}(X \times \mathbb{S}^1; G') \neq 0$ for some group G' . Because $\dim X \times \mathbb{S}^1 = n + 1$, $H_{n+1}(X \times \mathbb{S}^1; G') \cong \widehat{H}_{n+1}(X \times \mathbb{S}^1; G')$ and $H^{n+2}(\dim X \times \mathbb{S}^1, \mathbb{Z}) = 0$. Therefore, the exact sequence

$$\text{Ext}(H^{n+2}(X \times \mathbb{S}^1), G') \rightarrow \widehat{H}_{n+1}(X \times \mathbb{S}^1; G') \rightarrow \text{Hom}(H^{n+1}(X \times \mathbb{S}^1), G'),$$

where the coefficient group \mathbb{Z} in $H^{n+2}(X \times \mathbb{S}^1)$ and $H^{n+1}(X \times \mathbb{S}^1)$ are suppressed, yields that $H^{n+1}(X \times \mathbb{S}^1; \mathbb{Z}) \neq 0$. On the other hand, the Künneth formula from the proof of Theorem 3.1 (with \mathbb{Z} being the coefficient group in all cohomology groups) implies $H^{n+1}(X \times \mathbb{S}^1; \mathbb{Z}) = 0$, a contradiction. \square

Corollary 3.3. *Suppose for all $n \geq 1$ and all $X \in \mathcal{H}(n)$ the following holds: For every closed separator F of X with $\dim F = n-1$ there exists a group G such that either $H^{n-1}(F; G) \neq 0$ or $H_{n-1}(F; G) \neq 0$. Then there is no homogeneous metric AR -compactum Y with $\dim Y < \infty$.*

If $\mathcal{H}(G, n)$ denotes the class of all homogeneous metric ANR -compacta X with $\dim_G X = n$, the arguments from Theorem 3.1 provide the following result:

Proposition 3.4. *The following conditions are equivalent:*

- (1) $H^n(X; G) \neq 0$ for all $X \in \mathcal{H}(G, n)$ and all $n \geq 1$;
- (2) If $X \in \mathcal{H}(G, n)$ and $n \geq 1$, then $H^{n-1}(F; G) \neq 0$ for every closed set $F \subset X$ separating X ;
- (3) If $X \in \mathcal{H}(G, n)$ and $n \geq 1$, $H^{n-1}(F; G) \neq 0$ for every closed set $F \subset X$ separating X with $\dim_G F = n - 1$.

The corresponding homological analogue of Proposition 3.4 also holds for some groups G .

Proposition 3.5. *The following conditions are equivalent, where G is either a field or a torsion free group:*

- (1) $H_n(X; G) \neq 0$ for all $X \in \mathcal{H}(n)$ and all $n \geq 1$;
- (2) If $X \in \mathcal{H}(n)$, $n \geq 1$, and $F \subset X$ is a closed set separating X , then $H_{n-1}(F; G) \neq 0$;
- (3) If $X \in \mathcal{H}(n)$, $n \geq 1$, and $F \subset X$ is a closed set separating X with $\dim F = n - 1$, then $H_{n-1}(F; G) \neq 0$.

Proof. All implications except $(3) \Rightarrow (1)$ follow from the proof of Theorem 3.2. To prove $(3) \Rightarrow (1)$, we suppose there exists a space $X \in \mathcal{H}(n)$ with $H_n(X; G) = 0$. Considering the $(n + 1)$ -dimensional separator $X \times \mathbb{S}^1$ of $X \times \mathbb{S}^2$, we conclude that $H_{n+1}(X \times \mathbb{S}^1; G) \neq 0$. Because X and $X \times \mathbb{S}^1$ are ANR 's, their Čech homology groups are isomorphic

to the singular homology groups. Thus, we can apply the Künneth formula

$$\sum_{i+j=n+1} H_i(X) \otimes H_j(\mathbb{S}^1) \rightarrow H_{n+1}(X \times \mathbb{S}^1) \rightarrow \sum_{i+j=n} H_i(X) * H_j(\mathbb{S}^1),$$

where G is the coefficient group. Since $H_n(X; G) = H_{n+1}(X; G) = 0$ and $H_j(\mathbb{S}^1; G) = 0$ for all $j > 1$, $\sum_{i+j=n+1} H_i(X; G) \otimes H_j(\mathbb{S}^1; G) = 0$. If G is a torsion free group, then the group $\sum_{i+j=n} H_i(X; G) * H_j(\mathbb{S}^1; G)$ is also trivial because $H_1(\mathbb{S}^1; G) = G$ yields $H_{n-1}(X; G) * H_1(\mathbb{S}^1; G) = 0$. Therefore, $H_{n+1}(X \times \mathbb{S}^1; G) = 0$, a contradiction.

When G is a field, the group $H_{n+1}(X \times \mathbb{S}^1; G)$ is isomorphic to the trivial group $\sum_{i+j=n+1} H_i(X; G) \otimes H_j(\mathbb{S}^1; G)$. So, we have again a contradiction. \square

REFERENCES

- [1] P. Alexandroff, *Introduction to homological dimension theory and general combinatorial topology*, Nauka, Moscow, 1975 (in Russian).
- [2] R. H. Bing and K. Borsuk, *Some remarks concerning topological homogeneous spaces*, Ann. of Math. **81** (1965), no. 1, 100–111.
- [3] G. Bredon, *Sheaf Theory*, Second Edition, Graduate texts in Mathematics 170, Springer, 1997.
- [4] E. Effros, *Transformation groups and C^* -algebras*, Ann. of Math. **81** (1965), 38–55.
- [5] W. Massey, *Homology and cohomology theory*, New York, Dekker, 1978.
- [6] D. Montgomery, *Locally homogeneous spaces*, Ann. Math. **52** (1950), no. 2, 261–271.
- [7] E. Sklyarenko, *Homology theory and the exactness axiom*, Uspehi Mat. Nauk **24** (1969) no. 5(149), 87–140 (in Russian).
- [8] N. Steendrod, *Regular cycles of compact metric spaces*, Ann. of Math. **41** (1940), 833–851.
- [9] V. Todorov and V. Valov, *Alexandroff type manifolds and homology manifolds*, Houston J. Math., **40** (2014), no. 4, 1325–1346.
- [10] V. Valov, *Homogeneous ANR spaces and Alexandroff manifolds*, Topol. Appl. **173** (2014), 227–233.
- [11] V. Valov, *Local cohomological properties of homogeneous ANR compacta*, Fund. Math. **223** (2016), 257–270.

DEPARTMENT OF COMPUTER SCIENCE AND MATHEMATICS, NIPISSING UNIVERSITY, 100 COLLEGE DRIVE, P.O. BOX 5002, NORTH BAY, ON, P1B 8L7, CANADA

E-mail address: veskov@nipissingu.ca